

FULBRIGHT & JAWORSKI L.L.P.

A REGISTERED LIMITED LIABILITY PARTNERSHIP

600 CONGRESS AVENUE, SUITE 2400

AUSTIN, TEXAS 78701

WWW.FULBRIGHT.COM

TELEPHONE: 512.474.5201

FACSIMILE: 512.536.4598

Date: Thursday, June 21, 2007 10:07:04 AM**No. of pages (including cover):** 32**Matter No.:** 10413852**TO**

Examiner Goldberg

PHONE NO.**FACSIMILE NO.** 1-571-273-0743**FROM**

David Parker

PHONE NO. 512-536-3055

Should recipient confirm, by telephone, receipt of this facsimile transmission? No

COMMENTS:

Re: 09/832,069

Per your request...

CAUTION - CONFIDENTIAL

THE INFORMATION CONTAINED IN THIS FACSIMILE IS CONFIDENTIAL AND MAY ALSO CONTAIN PRIVILEGED ATTORNEY-CLIENT INFORMATION OR WORK PRODUCT. THE INFORMATION IS INTENDED ONLY FOR THE USE OF THE INDIVIDUAL OR ENTITY TO WHOM IT IS ADDRESSED. IF YOU ARE NOT THE INTENDED RECIPIENT, OR THE EMPLOYEE OR AGENT RESPONSIBLE TO DELIVER IT TO THE INTENDED RECIPIENT, YOU ARE HEREBY NOTIFIED THAT ANY USE, DISSEMINATION, DISTRIBUTION OR COPYING OF THIS COMMUNICATION IS STRICTLY PROHIBITED. IF YOU HAVE RECEIVED THE FACSIMILE IN ERROR, PLEASE IMMEDIATELY NOTIFY US BY TELEPHONE, AND RETURN THE ORIGINAL MESSAGE TO US AT THE ADDRESS ABOVE VIA THE U.S. POSTAL SERVICE. THANK YOU.

If you experience problems with this transmission or have questions, please call the sender.

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

APPLICANT: Runge, et al	§	ART UNIT:
	§	
FILED: April 10, 2001	§	
	§	
SERIAL NO.: 09/832,069	§	EXAMINER:
	§	
FOR: Mitochondrial DNA Damage as	§	
Predictor of Coronary	§	DOCKET:
Atherosclerotic Heart Disease	§	D6179CIP

Assistant Commissioner of Patents
BOX Patent Application
Washington, DC 20231

PRELIMINARY AMENDMENT UNDER 37 C.F.R. 1.111

Dear Sir:

Preliminary to the examination of the above-referenced application and in response to a Notice of Incomplete Reply mailed on September 21, 2001, Applicants respectfully request entry of the following amendments.

Attached hereto is a marked-up version of the changes made to the claims by the current amendment. The attached page is captioned "Version with markings to show changes made."

AMENDMENTS**IN THE SPECIFICATION:**

Please delete the paragraph beginning on page 14, line 20.

Please replace the paragraph beginning on page 15, line 13, with the following rewritten paragraph:

--Figures 7A and 7B show that decreased dietary protein intake correlates with less mitochondrial DNA damage in control and apoE mice. Control and apoE mice were fed either a 16% or 24% protein diet for 4 weeks, commencing at 6 weeks of age, and mitochondrial DNA damage was assessed by QPCR. Figure 7A illustrates the relative amplification of control and apoE mitochondrial DNA in aortas relative to the 24% protein fed control group. Less amplification product signifies increased mitochondrial DNA damage. Asterisks (*) indicate a significant difference between the 24% and 16% protein fed mice. Students' t' test P values are in parentheses. Figure 7B illustrates the relative amplification of control and apoE mitochondrial DNA in the left ventricle relative to the 24% protein fed control group. Less amplification product signifies increased mitochondrial DNA damage. Asterisks (*) indicate a significant difference between the 24% and 16% protein fed mice. Students' t' test P values are in parentheses.--

Please replace the paragraph beginning on page 16, line 8, with the following rewritten paragraph:

--Figure 8 shows that lipid peroxidation is significantly increased in apoE compared to control mice, but lipid peroxidation is significantly increased in control mice fed the western (21% fat) diet. Bar graph illustrates the level of lipid peroxidation in apoE and control mice fed chow (4% fat) or western (21%) diets. Values are expressed relative to the chow fed control mice. While apoE had significantly higher levels of lipid peroxidation compared to control mice regardless of diet, there were

no significant differences in the level of lipid peroxidation among the apoE fed the chow or western diets. By contrast, the control mice fed the western diet had significantly higher levels of lipid peroxidation (indicated by "**") compared to control mice fed the chow diet.--

Please replace the paragraph beginning on page 16, line 20, with the following rewritten paragraph:

--**Figure 9** shows QPCR of age-matched normal and atherosclerotic aortic mtDNA from humans. The 16.2 kb (upper row) and 0.22 kb (bottom row) products represent the long and short human QPCR amplicons, respectively. Methods for QPCR from human mtDNA have been previously described (Ballinger, 1996, 1999; Yakes, 1997).--

Please replace the paragraph beginning on page 17, line 5, with the following rewritten paragraph:

--**Figures 10A and 10B** show that hybrid apoE (-/-), SOD2 (+/-) mice have increased atherosclerotic lesions and mtDNA damage. In **Figure 10A**, whole aortas from hybrid apoE (-/-), SOD2 (+/-) and apoE (-/-) littermates were oil-red-O stained. The upper left panel shows an oil-red-O stained aorta from a 17 week-old apoE (-/-), SOD2 (+/-) male (4% fat diet), and the upper right panel shows the results from an apoE (-/-) male littermate (4% fat diet). Lower left and right panels show stained aortas from 34 week-old apoE (-/-), SOD2 (+/-) and apoE (-/-) littermates, respectively. Black arrows indicate presence of large atherosclerotic lesion in the apoE (-/-), SOD2 (+/-) and apoE (-/-), (not yet present in the 17 week-old apoE (-/-) littermate), and the red arrow in each panel point to an example of smaller lesion formed at arterial branches, that were more frequent and characteristics of the apoE (-/-), SOD2 (+/-) mice. **Figure**

10B is a bar graph depicting atherosclerotic lesion frequencies from apoE (/), SOD2 (+/-) and apoE (-/-) mice whole aortas, visualized by oil-red-O staining (n=4).--

Please replace the paragraph beginning on page 18, line 1, with the following rewritten paragraph:

--Figure 11 shows the increased incidence of mitochondrial DNA damage in patients with risk factors for myocardial infarction. Blood samples were taken at the time of cardiac catheterization to measure mitochondrial DNA damage by QPCR. For this figure, values of mitochondrial DNA damage above the mean for a "normal" population were considered increased. The bar graphs show the percentage of patients in each category with increased mitochondrial DNA damage. The occurrence of increased mitochondrial DNA damage is more frequent in subjects with hypertension, cigarette smoking, diabetes and combinations of these risk factors for myocardial infarction.--

Please replace the paragraph beginning on page 18, line 12, with the following rewritten paragraph:

--Figure 12 shows mitochondrial DNA damage before and after a 20 mile "training run". Blood samples were taken, a buffy coat was obtained and mitochondrial DNA damage was assessed by QPCR. Although mitochondrial DNA damage is present immediately after the training run, this quickly returns to normal values as shown here.--

Please replace the paragraph beginning on page 18, line 18, with the following rewritten paragraph:

--Figure 13 shows that ultramarathoners had significantly less mitochondrial DNA damage after being fed a high fat meal than did the control subjects. In all cases, mitochondrial DNA damage was assessed before, immediately after eating a high fat meal and 4-6 hours later. The observed mitochondrial DNA damage in controls reflects the immediate effects of dietary intake of ROS in the form of fats and cholesterol. The relative "protection" observed in the ultramarathoners may relate to an upregulation of anti-oxidant defense mechanisms in these individuals.--

Please replace the paragraph beginning on page 19, line 6, with the following rewritten paragraph:

--Figure 14 shows that ultramarathoners had significantly less protein damage after being fed a high fat meal than did the control subjects--

Please replace the paragraph beginning on page 53, line 3, with the following rewritten paragraph:

--Aortic tissues from the hypercholesteremic apoE mice fed the chow diet had significantly increased levels of mitochondrial DNA damage compared to controls ($P < 0.05$; data not shown). These differences were observed in both the 10 and 34 week old mice, which clearly showed that apoE aortas sustained greater mitochondrial DNA damage *in vivo* compared to "healthy" mice. The 10 week old apoE aorta yielded a 61% decrease in amplification (decreased amplification correlates with increased DNA damage) relative to the zero class control, or an estimated 0.582 ± 0.123 mitochondrial DNA lesions/10 kilobases (kb) compared to 0.0 ± 0.198 lesions/10 kb in the 10 week old control mouse ($P=0.018$). Similarly, the 34 week old apoE group had a three-fold increase in estimated

mitochondrial DNA damage compared to the age matched control group (1.325 ± 0.257 lesions/10 kb versus 0.453 ± 0.162 lesions/10 kb; $P=0.007$). These same patterns were also observed in tissue of the left ventricle of the heart. (data not shown). For example, the 10 week old apoE mice had a 67% decrease in amplification compared to the zero class control (0.685 ± 0.093 lesions/10 kb versus 0.0 ± 0.048 lesions/10 kb; $P < 0.001$), while the 34 week old apoE mice sustained approximately 4-fold higher estimated mitochondrial DNA lesions compared to the age-matched controls (0.819 ± 0.151 lesions/10 kb versus 0.213 ± 0.295 lesions/10 kb, $P = 0.056$).--

Please replace the paragraph beginning on page 54, line 4, with the following rewritten paragraph:

--The effects of the western diet were also assessed in apoE and control mice. In the control mice, the western diet was associated with higher levels of mitochondrial DNA damage in the 10 week-old control group aortas compared to chow fed control mice (0.31 ± 0.17 lesions/10 kb versus 0.0 ± 0.20 lesions/10 kb, western and chow, respectively), however, this difference was less significant ($P = 0.25$). Such differences were less obvious in the 34 week old control mice (i.e. 0.45 ± 0.16 lesions/10 kb and 0.49 ± 0.25 lesions/10 kb in chow versus western, respectively, $P = 0.90$). Similarly, the 10 week old control mice showed increased estimated mitochondrial DNA damage with the western diet in heart (58% decrease in amplification, 0.54 ± 0.23 lesions/10 kb and 0.0 ± 0.05 lesions/10 kb in western versus chow, respectively), yet similar with the results seen in the aorta, no obvious differences were seen in the 34 week old control mice (0.21 ± 0.29 lesions/10 kb and 0.25 ± 0.21 lesions/kb in chow versus western, respectively). By contrast, no apparent increases in mitochondrial

DNA damage accompanied the western diet in either aorta or heart tissues from apoE mice. Hence, increased dietary fat appeared to be associated with increased mitochondrial DNA damage in the 10 week old control mice only.--

Please replace the paragraph beginning on page 55, line 20, with the following rewritten paragraph:

--Because no significant differences in mitochondrial DNA damage were observed between the chow and western diets with matched genotype and age, the effects of protein in the 10 week old mice were investigated. Both apoE and c57Bl6 control mice were fed either 16% or 24% protein diets for 4 weeks, commencing at 6 weeks of age. While both diets resulted in apoE still having greater levels of mitochondrial DNA damage compared to controls (aorta $P = 0.015$; heart $P = 0.005$), the lower protein diet resulted in a significant decrease in damage in both control ($P = 0.007$) and apoE ($P < 0.001$) in aorta (Figure 7). This same trend was also observed in apoE heart ($P = 0.002$). Hence, the lower protein diet was associated with decreased mitochondrial DNA damage in both types of mice.--

Please replace the paragraph beginning on page 58, line 8, with the following rewritten paragraph:

--Lipid peroxidation levels were measured by determining the levels of MDA and 4-HNE. The hypercholesteremic apoE mice had significantly increased levels of lipid peroxides compared to control mice ($P < 0.05$). However, in contrast with the results observed with the total cholesterol levels, the degree of lipid peroxidation did not significantly increase in apoE mice on the western compared to the chow diet (Figure 8).

By contrast, the control mice fed the western diet sustained increased lipid peroxidation compared to the chow diet ($P < 0.05$). Hence, while the apoE mice always had the highest levels of lipid peroxides compared to the control mice, the degree of lipid peroxidation in apoE did not increase when fed the western diet compared to the chow diet.--

Please replace the paragraph beginning on page 60, line 4, with the following rewritten paragraph:

--QPCR analysis revealed significantly increased mtDNA damage in atherosclerotic tissues compared with healthy control group aortas (Figures 9 and Table 8). By contrast, a marker for nuclear DNA damage, the *β -globin* gene cluster, did not sustain significant damage in atherosclerotic aorta compared with controls ($P = 0.15$). Consequently, the mtDNA sustained significant and preferential damage in human atherosclerotic aortas. These findings were consistent with the notion that mtDNA damage is an integral feature of atherogenesis. Because increased mtDNA damage was a consistent feature in both human and mouse atherogenesis, it was hypothesized that mitochondrial dysfunction is an important event in atherogenesis, possibly modulating the phenotype of vascular cells that are integral to atherosclerotic lesion formation.--

Please replace the paragraph beginning on page 62, line 11, with the following rewritten paragraph:

--For analysis of atherogenesis and mtDNA damage, apoE (-/-), SOD2 (+/-) mice and apoE (-/-) controls were fed a 4% fat diet and sacrificed at 17 weeks of age. Assessment of functional SOD2 activity in hybrid apoE (-/-), SOD2 (+/-) aortas demonstrated a 44% reduction of activity compared with "normal" apoE (-/-) mice (data not shown; $P <$

0.001). Histological examination of atherosclerotic lesion development in these mice revealed that apoE (-/-), SOD2 (+/-) mice developed earlier lesions at physiologically relevant sites, including in the ascending aorta at the origin of the cerebral vessels. However, the most impressive finding was that at 17 weeks of age, oil-red-O positive lesions were present throughout the descending aorta at virtually every branch point of the spinal arteries in the apoE (-/-), SOD2 (+/-) mice (Figure 10A). Analogous lesions were rare in both 17-week-old and 34-week-old apoE (-/-) mice. Overall, the apoE (-/-), SOD2 (+/-) mice had 2.5 fold as many atherosclerotic lesions ($P = 0.02$) compared with apoE (-/-) littermates (Figure 10B). Although the small size of atherosclerotic lesions at the branch points precluded direct measurement of mtDNA damage at these sites, quantification of aortic mtDNA damage through QPCR revealed increased aortic mtDNA damage in the apoE (-/-), SOD2 (+/-) mice compared with their age-matched apoE (-/-) littermates (Table 8; $P = 0.006$). Thus, while providing further evidence that mitochondrial dysfunction, resulting from oxidative mtDNA damage is important in atherogenesis, these findings also supports the notion that, under normal circumstances, decreased expression of SOD2 in response to turbulent flow at arterial branch points promotes atherogenesis at these sites.--

Please replace the paragraph beginning on page 77, line 9, with the following rewritten paragraph:

--Patient who were undergoing cardiac catheterization were recruited for an *in vivo* study of mitochondrial DNA damage. All signed informed consent for an additional blood sample to be taken at the time of cardiac catheterization. In all, there were 75 patient. White blood cells were isolated by standard techniques (the "buffy coat" preparation), and DNA

(both nuclear and mitochondrial) was isolated. Mitochondrial DNA damage was determined by quantitative PCR using both "short mitochondrial DNA fragments" and damage within the β -globin gene as controls. Figure 11 shows the increased incidence of mitochondrial DNA damage in patients grouped according to risk factors associated with myocardial infarction.--

Please replace the paragraph beginning on page 78, line 18, with the following rewritten paragraph:

--To examine further the extent of mitochondrial DNA damage under a variety of *in vivo* conditions, 5 individuals who are "ultramarathoners" and an age-matched group of 6 individuals who are healthy and active, but not ultramarathoners, were examined. First, mitochondrial DNA damage was measured before and after a 20 mile "training run" by the ultramarathoners. Blood samples were taken, a buffy coat was obtained, and mitochondrial DNA damage was assessed. Figure 12 shows that there is a mild increase in mitochondrial DNA damage just after the run, which returns to normal within 4 hours. On a different day, the ultramarathoners and the control subjects were fed a high fat meal. The high fat meal comprised the "El Grande Platter" at a local Mexican food restaurant, which contained approximately 3000 calories, of which approximately 60-70% were fat calories. Blood samples were taken just before the meal, and 2 and 6 hours after. Buffy coats and DNA isolation were performed. Mitochondrial protein damage was also quantitated using white blood cell mitochondrial protein as a source. Figures 13 and 14 show that there is a gradual, small increase in mitochondrial DNA damage in the controls, and increased mitochondrial protein damage at 2 hours that decreases in 4 hours. In the ultramarathoners, mitochondrial DNA damage and protein damage actually


decreased to below baseline levels at 2 hours and begins returning to baseline by 6 hours. Presumably, since the ultramarathoners are subjected to increased oxidative stress when they run, their anti-oxidant protective systems are more effective. Thus, when exposed to a stimulus for DNA damage, the ultramarathoners have even less damage than normal. Subsequently, their antioxidant systems are activated and decrease their oxidant levels to below resting levels.--

REMARKS

Applicants submit that these Amendments delete all reference to Figures 7A and 7B and incorporate the pertinent information regarding Figures 7A and 7B from the Brief Description of Drawings into the Specification. No new matter is contained in these amendments.

Respectfully submitted,

Date: Nov 21, 2001
ADLER & ASSOCIATES
8011 Candle Lane
Houston, Texas 77071
(713) 270-5391
badler1@houston.rr.com


Benjamin Aaron Adler, Ph.D., J.D.
Counsel for Applicant
Registration No. 35,423